INFOMAR Integrated Mapping for the Sustainable Development of Ireland's Marine Resource
An Roinn Aeráide, Fuinnimh agus Comhshaoil Department of Climate, Energy and the Environment
Geological Survey Suirbhéireacht Gheolaíochta Ireland | Éireann
Marine Institute Foras na Mara



Marine Geoscience Report 001

Waterford Harbour & Shipping Channel

Overview

Waterford Harbour and its shipping channel form a funnel-shaped estuary in south-east Ireland, linking the Port of Waterford to the Celtic Sea. Spanning the counties of Waterford, Wexford and Kilkenny, the channel has long supported navigation, trade and settlement in the region.



Figure 1: Bathymetry of Waterford Estuary and shipping channel. Depths are referenced to Lowest Astronomical Tide (LAT).

The estuary developed during the Quaternary Period through river erosion, glaciation and later marine flooding. Glacial advances eroded valleys and deposited sediment. As sea level rose, tidal currents and waves reworked these sediments, helping to shape the modern estuary and shipping channel. The underlying Devonian sandstones and Carboniferous limestones, intersected by faults and fractures, helped define the estuary's boundaries and influenced the alignment of river channels. Tidal currents, river input and sediment transport continue to shape the harbour through erosion and deposition.

History

Waterford's strategic geographic position has made Waterford Harbour one of the main gateways to Ireland since prehistoric times, with evidence of early settlements found along the shores. The Vikings established the port town in 914 AD, naming it *Vedrarfjordr*, often interpreted as "haven from windy seas", which evolved into Waterford over time (Foley, 2021).




Figure 2: Nautical Chart of Waterford Harbour 1787. Image from (Wiki Commons, 2022).

The region has a long maritime history, including boatbuilding, fishing, lighthouses, forts, weirs, shipwrecks, and leisure activities. Historical port infrastructure developed during the medieval period expanded significantly in the 18th and 19th centuries, aligning with the growth of industrial shipping (Foley, 2021).

The harbour also provides access to the Suir, Barrow and Nore river systems. At the confluence of the Rivers Suir and Barrow, INFOMAR bathymetry data show deeper channels along outer bends and sediment banks along inner bends (Figure 3). Stronger currents scour the outer bends, while slower flow along the inner bends allows sediment to accumulate. Differential erosion also influences channel morphology, as softer sediments are more readily eroded, while more resistant materials remain and help shape the channel.




Figure 3: Bathymetry at the meeting of the Rivers Suir and Barrow with a modern nautical chart background.

Extensive dredging operations were conducted in the 20th century to maintain navigability for larger vessels. The bathymetry data shows the probable spoil heap (dredge dump site) at the mouth of the harbour, rising 4 m above the surrounding seabed. While dredging interventions enhanced commercial capacity, they also altered natural sedimentary and hydrological regimes. Current management aims to balance economic utility with ecological preservation, as well as monitoring the spoil heap to ensure pollutants do not spread (Port of Waterford, 2023).




Figure 4: Probable spoil heap at the mouth of the harbour, rising to over 4 m above the surrounding seabed.

Another notable feature in the harbour is the shipwreck of the SS Harvard. In 1870, while en route from Spain to Glasgow, the vessel struck rocks in Waterford Harbour (INFOMAR, 2013). The wreck lies near the northern bank of the river, at a depth of approximately 12 m, as shown in Figure 5.




Figure 5: Point cloud imagery of the SS Harvard on the seabed, with overview location to the left.

Regional Geology

The geological setting of the Waterford Shipping Channel reflects the long development of south-east Ireland's landscape. Its bedrock foundation was shaped by sedimentation, mountain-building and faulting over hundreds of millions of years, while more recent fluvial, glacial and marine processes modified the estuary and seabed.




Figure 6: Geological map of south-east Ireland.

The region's bedrock is predominantly composed of Ordovician shales to the east of Waterford and Devonian Old Red Sandstone (ORS) to the west. The Ordovician shales formed in the marine basin of the Iapetus Ocean during the Ordovician (485 – 444 Ma), while the Devonian ORS was deposited in a largely terrestrial environment during the Devonian (419 – 359 Ma), with sediment supplied from mountain ranges to the north (Tietzsch-Tyler, et al., 1994). These are interspersed with Silurian (444 – 419 Ma) volcanic rocks, formed as the Iapetus Ocean closed, creating a varied geological landscape.

During the Carboniferous Period (359 – 299 Ma), warm tropical seas flooded low-lying areas, leading to the deposition of fossil-rich limestone across much of south-east Ireland. At the end of the Carboniferous, during the Variscan Orogeny around 300 Ma, tectonic forces folded the Devonian and Carboniferous rocks into broad E – W structures. Subsequent erosion exposed Devonian sandstones in the uplands, while limestone formed the bedrock in the lower-lying parts of the folds (Parkes, et al., 2012).




Figure 7: Geological map of Waterford Estuary.

Structural geology has strongly influenced Waterford Harbour. Faults and fractures in the underlying bedrock helped guide the alignment of the estuary and river channels, and contributed to the formation of the harbour's boundaries. These controls continue to influence channel form, sediment transport and patterns of erosion and deposition.

More recently, alluvium and marine sediments accumulated in estuarine zones. The channel substrate is composed of a mix of silt, sand and gravel, as shown in Figure 8, and these sediments are continually redistributed by tidal forces and river input. This sediment movement influences the harbour's physical conditions and habitats (Parkes, et al., 2012).




Figure 8: Sediment classification map of Waterford Harbour showing the distribution of muds, sands and coarse sediments.

Around Little Island, the seabed morphology shown in Figure 9 highlights variations in sediment deposition and erosion. The southern flank has more pronounced bends, where deeper scour occurs along outer curves, while sediment banks accumulate nearer the shore. This suggests that stronger currents accelerate through the bends and erode the channel floor, while lower-energy areas promote deposition. These patterns reflect the interaction between tidal currents, river flow and channel morphology.




Figure 9: Bathymetry around Little Island.

Biodiversity

The Waterford Shipping Channel supports biodiversity within its intertidal, estuarine and riverine habitats. The channel is associated with the Lower River Suir Special Area of Conservation (SAC) and the River Barrow and River Nore SAC, which include tidal reaches, estuarine habitats and species such as Atlantic Salmon, Sea Lamprey, River Lamprey, Twaite Shad and Otter (National Parks & Wildlife Service, 2024) (National Parks & Wildlife Service, 2023). Mudflats and saltmarshes support salt-tolerant plants such as glasswort and sea aster, while birds including Oystercatcher and Curlew use intertidal areas for feeding and roosting (National Biodiversity Data Centre, 2017).



Figure 10: A curlew on the coast of Waterford. Image from (Bird Watching Ireland, n.d.).

The harbour also supports migratory aquatic species, including Atlantic Salmon and European Eel, which use the river system as a migratory pathway. The variety of mudflats, saltmarshes, channels, beaches and rocky shores provides a range of habitats, but these environments are sensitive to coastal development, pollution, dredging and changes in sediment supply (National Parks & Wildlife Service, 2024) (National Parks & Wildlife Service, 2023). Conservation designations under the EU Habitats Directive help recognise and manage the ecological value of the wider estuary (National Parks and Wildlife Service, 2019).

Conclusion

Waterford Harbour's shipping channel reflects the combined influence of bedrock structure, river flow, glacial modification, dredging and ongoing tidal reworking. Its funnel-shaped estuary provides naturally sheltered deep water, while sediment transport, erosion and deposition continue to shape the channel. The harbour's long use for settlement, trade and navigation reflects these physical advantages. Understanding the balance between geology, sediment movement, human intervention and ecological value is important for the future management of the channel.

Bibliography

- Bird Watching Ireland, n.d. *Birding-Dungarvan-Waterford.* [Online]
  Available at: https://www.birdwatchingireland.com/birding-dungarvan-waterford#bird-watching-royal-canal-binns-bridge-6029-8

- Foley, N. S., 2021. *The Maritime Heritage of Waterford Estuary,* s.l.: Port of Waterford.

- INFOMAR, 2013. *SS Harvard wreck Information Sheet,* s.l.: s.n.

- National Biodiversity Data Centre, 2017. *Irish Vegetation Classification.* [Online]
  Available at: https://biodiversityireland.ie/app/uploads/2017/01/SM1B.pdf

- National Parks & Wildlife Service, 2023. *Site Synopsis: Lower River Suir SAC,* s.l.: Department of Housing, Local Government and Heritage.

- National Parks & Wildlife Service, 2024. *Site Synopsis: River Barrow and River Nore SAC,* s.l.: Department of Housing, Local Government and Heritage.

- National Parks and Wildlife Service, 2019. *The Status of EU Protected Habitats and Species in Ireland,* s.l.: Department of Culture, Heritage and the Gaeltacht.

- Parkes, M., Meehan, R. & Préteseille, S., 2012. *The Geological Heritage of Waterford: An Audit of County Geological Sites in Waterford,* Dublin: Irish Geological Heritage Programme, Geological Survey Ireland.

- Port of Waterford, 2023. *Port of Waterford PERS Environmental Report,* s.l.: s.n.

- Tietzsch-Tyler, D. et al., 1994. *Geology of South Wexford,* s.l.: Geological Survey Ireland.

- Wiki Commons, 2022. *An actual survey of the harbour and river of Waterford and of the bay of Tramore.* [Online]
  Available at:
  https://commons.m.wikimedia.org/wiki/File:An_actual_survey_of_the_harbour_and_river_of_Waterfor_and_of_the_bay_of_Tramore..._-_btv1b53011020p_%281_of_2%29.jpg